

Explaining the HERA Anomaly Without Giving Up R-parity Conservation

B. Dutta¹, R. N. Mohapatra² and S. Nandi³

⁽¹⁾ *Institute of Theoretical Science, University of Oregon, Eugene, OR 97403*

⁽²⁾ *Department of Physics, University of Maryland, College Park, Md-20742, USA.*

⁽³⁾ *Department of Physics, Oklahoma State University, Stillwater, OK 74078*

(April, 1997)

We point out that in extended supersymmetric models such as supersymmetric left-right models, it is possible to have leptoquarks that explain the HERA high Q^2 anomaly without giving up R-parity conservation. The leptoquarks belong to vectorlike $(2, 2, \pm \frac{4}{3}, 3 \text{ or } 3^*)$ representations of $SU(2)_L \times SU(2)_R \times U(1)_{B-L} \times SU(3)_c$ (denoted by G_{2213}). Considerations of D-terms imply that the only phenomenologically viable model is the one where the leptoquark couples to positrons and the up quark. Unlike the R-parity violating scenario, the leptoquarks are accompanied by new superpartners, the leptoquarkino which leads to many interesting signatures in other collider experiments. At Tevatron, pair productions of the leptoquarkino will give rise to dilepton signals very distinct from the top productions. These models can lead to unification of gauge coupling constants at a scale of around 10^{10} GeV implying that grand unification group is not of the usual $SU(5)$ or $SO(10)$ types but rather an automatically R-parity conserving $SU(5) \times SU(5)$ GUT model recently proposed by one of the authors (R. N. M.) which leads to a stable proton.

OITS-627

UMD-PP-97-108

OSURN-325

If the high Q^2 anomaly observed recently in the e^+p scattering by the H1 [1] and the ZEUS [2] collaborations is confirmed by future data, it will be an extremely interesting signal of new physics beyond the standard model. A very plausible and widely discussed interpretation of this anomaly appears to be in terms of new scalar particles capable of coupling to e^+u or e^+d of scalar leptoquarks [3] with mass around 200 GeV. Alternative interpretations based on contact interactions [4] or a second Z' [5] have been proposed; but attempts to construct models that lead to the desired properties seem to run into theoretical problems.

The leptoquark must be a spin zero color triplet particle with electric charge $5/3$ or $2/3$. The latter electric charge assignment makes it possible to give a plausible interpretation of the leptoquark as being the superpartner of the up-like quark [6] of the minimal supersymmetric standard model (MSSM) provided one includes the R-parity violating couplings of the type $\lambda' QLd^c$ to the MSSM (where Q, L denote the quark and the lepton $SU(2)_L$ doublets and u^c, d^c, e^c denote the $SU(2)_L$ singlets [7]). There are however very stringent upper limits on several R-parity violating couplings: for instance, if in addition to the λ' term described above, one adds the allowed $\lambda'' u^c d^c d^c$ terms to the superpotential, then it leads to catastrophic proton decay unless $\lambda' \lambda'' \leq 10^{-24}$ [8]. There are also stringent limits on $\lambda'_{111} \leq 10^{-4}$ [9] from neutrinoless double beta decay, which forces the leptoquark to be $\tilde{c}$ or $\tilde{t}$ rather than the obvious choice $\tilde{u}$. Furthermore, within such a framework, the lightest supersymmetric particle (LSP) is no more stable and therefore, there is no cold dark matter (CDM) candidate in such theories. In view of the fact providing a natural

CDM candidate was long considered an attractive feature of the supersymmetric models, it may be worthwhile to consider extensions of the MSSM which incorporate the leptoquark without giving up R-parity conservation (and hence the idea of LSP as a natural CDM candidate). It is the goal of this letter to report on the results of such a study.

The class of supersymmetric theories where R-parity conservation is automatic can provide an absolutely stable LSP that can act as the CDM candidate. We will therefore use them as a typical framework for studying the consequences of leptoquarks incorporated into such theories while at the same time maintaining automatic R-parity conservation. Minimal versions of such theories are based on the gauge group $SU(2)_L \times U(1)_{I_{3R}} \times U(1)_{B-L}$ or $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ [10] with usual assignments for the quarks and leptons. We will consider the latter left-right symmetric gauge group. The quark doublets $Q \equiv (u, d)$ and $Q^c \equiv (u^c, d^c)$ transform as doublets under the $SU(2)_L$ and $SU(2)_R$ groups respectively and similarly the lepton doublets $L \equiv (\nu, e)$ and $L^c \equiv (\nu^c, e^c)$ respectively. The $SU(2)_R$ gauge symmetry breaking is achieved via the $B-L$ non-singlet isotriplets $\Delta^c \equiv (1, 3, -2)$ and $\bar{\Delta}^c \equiv (1, 3, +2)$ and their lefthanded counterparts added to maintain left-right symmetry (the numbers in the parenthesis denote the $SU(2)_L, SU(2)_R$ and $U(1)_{B-L}$ quantum numbers). The standard model symmetry is broken by the bi-doublet fields $\phi(2, 2, 0)$.

We augment this model by including the leptoquark fields. Demanding that the lepto-quarks couple to e^+d or e^+u leads to the conclusion that they must belong to the multiplet $(2, 2, 4/3, 3)$ (denoted Σ_{QL^c}) and $(2, 2, -4/3, 3^*)$ (denoted by $\bar{\Sigma}_{Q^cL}$). Each of these multi-

plets have four scalar fields and four fermion fields which will be denoted in what follows by the obvious subscript corresponding to their couplings. We denote the four scalar leptoquarks as $\bar{\Sigma}_{ue^c}$, $\bar{\Sigma}_{uv^c}$, $\bar{\Sigma}_{de^c}$ and $\bar{\Sigma}_{d\nu^c}$ and their fermionic partners (to be denoted by a tilde on the corresponding scalar field). Writing down the most general superpotential, one can easily convince oneself that the resulting theory maintains the property of automatic R-parity conservation. Thus the lightest neutralino LSP will be stable in this model and can serve as the cold dark matter.

Before discussing the application of this model to explain the HERA anomaly, let us first discuss the mass spectrum of the model. The superpotential for this model will have a direct mass term of the form $M_0 \Sigma \bar{\Sigma}$ which will imply that the fermionic fields in the leptoquark multiplet will have a masses M_0 prior to symmetry breaking. There may be other contributions to these masses from radiative corrections which will split the degeneracy implied by the above mass term.

As far as the scalar leptoquark states are concerned, their masses will receive several contributions: first a direct common contribution from the M_0 term given above. After symmetry breaking, the D-terms of the various gauge groups will contribute. There is also soft SUSY breaking contribution along with the radiative correction. Assuming that the $SU(2)_R$ symmetry is broken by the vev's $\langle \Delta^c \rangle = v_1$, $\langle \bar{\Delta}^c \rangle = v_2$ and the $SU(2)_L$ symmetry is broken by the two ϕ vev's as $\text{diag} \langle \phi_u \rangle = (0, v \sin \beta)$ and $\text{diag} \langle \phi_d \rangle = (v \cos \beta, 0)$, we can write the masses for the various scalar leptoquarks as follows:

$$M_{\Sigma_a}^2 = M_0^2 + (I_{R_a}^2 g_{2R}^2 - \frac{B-L}{8} g_{B-L}^2)(v_1^2 - v_2^2) \quad (1)$$

$$+ \frac{g_{2L}^2 v^2 I_{3L}}{4} \cos 2\beta + \Delta_m^2 + \text{Radiative correction}$$

Δ_m^2 is the soft SUSY breaking contribution. The values of the $I_{R_a}^2$ and $B-L$ are given in Table I:

states	$I_{R_a}^2$	$B-L$	I_{3L}
$u^c e$	$\frac{1}{2}$	$\frac{4}{3}$	$\frac{1}{2}$
$u^c \nu$	$\frac{1}{2}$	$\frac{4}{3}$	$-\frac{1}{2}$
$u e^c$	$-\frac{1}{2}$	$-\frac{4}{3}$	$-\frac{1}{2}$
$d e^c$	$-\frac{1}{2}$	$-\frac{4}{3}$	$\frac{1}{2}$
$d^c \nu$	$-\frac{1}{2}$	$\frac{4}{3}$	$-\frac{1}{2}$
$d^c e$	$-\frac{1}{2}$	$\frac{4}{3}$	$-\frac{1}{2}$

TABLE I. The I_{3R} and B_L quantum numbers of the various leptoquark states

The radiative correction can be positive or negative. From table I and Eq.(1), we see easily that if $v_1 < v_2$, then the lightest leptoquark state is the first entry in the table I which corresponds to the conjugate of $\bar{\Sigma}_{ue^c}$ since $g_{2R}^2 > 2g_{B-L}^2$ in the left-right models. Furthermore, interestingly enough for this choice of vev's, assuming the combined $\Delta_m^2 + \text{Radiative correction}$ to be smaller than the D-term, the leptoquarkino states are heavier than the lightest leptoquark state. We assume their masses to be in the range of 300 to 400 GeV. In a subsequent section we will obtain a lower bound on the leptoquarkino mass from the present collider data.

Turning to the couplings of the leptoquarks Σ and $\bar{\Sigma}$ to quarks and leptons, it is given by the superpotential:

$$W_{lq} = \lambda_{ij} (\Sigma Q_i^c L_j + \bar{\Sigma} Q_i L_j^c) \quad (2)$$

(where i, j are generation indices). Let us assume for simplicity that λ_{ij} are diagonal in the mass eigenstate basis for the quarks and leptons. Explanation of the HERA high Q^2 anomaly seems to require $\lambda_{11} \simeq 0.05$ which we will assume from now on and mass of the scalar field in Σ (assumed to be Σ_{ue^c} from the above mass arguments) around 200 GeV. As far as the other couplings go, they will be strongly constrained by the present experimental upper limits on the low energy processes such as $\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$ etc. which will arise at the one loop level from the exchange of Σ and $\bar{\Sigma}$. There are also tree level diagrams which can lead to rare processes such as $K \rightarrow \pi e^- \mu^+$. These processes imply an upper limit of $\lambda_{22} \leq 2 \times 10^{-3}$. The most stringent upper limit on λ_{22} comes from the upper limit on the process $K_L^0 \rightarrow \mu^+ e^-$ and yields $\lambda_{22} \leq 2 \times 10^{-4.5}$. Turning to λ_{33} , the most stringent limits arise from the present upper limit on the branching ratio for the process $\tau \rightarrow e\gamma$ which the 1996 Particle data tables give as $\leq 1.1 \times 10^{-4}$ [11]. This implies a weak upper limit on $\lambda_{33} \leq 0.2$. Thus the third generation leptoquark coupling λ_{33} could in principle be comparable to λ_{11} . If the efficiency for the detection of τ leptons at HERA were comparable to the detection efficiency for electrons, that could also severely limit λ_{33} . Finally we also note that such a value for λ_{11} is also consistent with the present data on the parity violation in atomic physics. Thus it appears that our leptoquark couplings are consistent with all known low energy data. It is clear from the above discussion that the leptoquarks do not respect $\mu - e$ universality.

The existence and properties of the leptoquarkino provides a new and unique signature for our model as compared to all other proposals to explain the HERA anomaly. To see this note that in hadron colliders, we can produce pairs of leptoquarkinos at the same rate as the $t\bar{t}$ pair due to identical color content. Moreover, due to R-parity conservation the leptoquarkino decay leads to a missing energy signal as follows: $\tilde{\Sigma} \rightarrow e^+ \tilde{u}$ with $\tilde{u} \rightarrow u + \chi^0$ or $\tilde{\Sigma} \rightarrow u + e^+$ with $e^+ \rightarrow e^+ + \chi^0$. In both the cases we have e^+, u plus missing energy in the final state. If the λ_{33} coupling is comparable to λ_{11} as is allowed, then the branching ratio for leptoquarkino decay

to electrons will be 50%. This signal is similar to the top signal at the Tevatron with one crucial difference. The dilepton branching ratio from the leptoquark pairs is 100% compared to only 10% from the top pairs. There will be no $\mu^+\mu^-$ events. The branching ratio to e^+e^- will be 25% compared to only 1% from the top. The present observations should therefore lead to lower limits on the mass of the lightest leptoquarkino pairs. Thus an excess of dilepton pairs over that expected from the top productions, or an excess in e^+e^- channel over $\mu^+\mu^-$ will be a clear signal of leptoquark productions at Tevatron. In addition, leptoquarks will also give rise to harder leptons and larger missing energy events. In Fig. 1, we plot the cross section for the leptoquarkino pair production. We see that for the present combined Tevatron data of about 200 pb^{-1} , there would be 6 events of type e^+e^-jj plus missing energy for $M_{\tilde{\Sigma}_{u^c e}} = 250 \text{ GeV}$, assuming a detection efficiency of 0.2 for the e^+e^- mode. At present CDF collaboration has only one such event [12], and the D0 collaboration has also 1 event [13]. Using these two events, and the cross sections given in Fig. 1, we obtain a lower bound on the leptoquarkino mass of about 290 GeV (assuming the detection efficiency to remain 0.2 for the higher masses). Any evidence for an excess of dielectron events would be interpretable in terms of this new leptoquarkino. We urge the CDF and D0 collaboration to look for such excess, and also to look for any dilepton event in which the lepton P_T or missing energy do not fit the top productions. The recent CDF analysis [14] shows that the leptoquark mass bound at 95% CL is 210 GeV assuming the branching ratio is 1. In this model however the branching ratio is less than 1, since the leptoquark can also decay into the top quark and the τ lepton.

So far we have discussed the signals of leptoquarkinos in the usual SUSY theories. If however the supersymmetry breaking is communicated to the observable sector by the SM gauge interaction [15], the signal is different. Let us first consider the case when neutralino is the NLSP in these models. The leptoquarkino decays into a spositron and u quark (squarks masses are large in these models). The spositron decays (100%) into a positron and a neutralino and the neutralino then decays (100%) into a hard photon and a gravitino. Consequently the final state in the leptoquarkino pair production process has $e^+e^-\gamma\gamma jj$ plus missing energy without any SM background. If lighter stau is the NLSP and neutralino is the NNLSP, spositron decays (100%) into a positron and a neutralino and the neutralino then decays (100%) into a τ and a lighter stau. The lighter stau then decays into a tau and a gravitino. Consequently the final state in the leptoquarkino pair production process in the Tevatron has $e^+e^-2\tau^+2\tau^-jj$ plus missing energy. Out of these six leptons in the final state, one τ pair (produced from the decay of stau) has much higher P_T than the other leptons.

One could also look for the signals of leptoquarkino

production in $e^+\gamma$ colliders. The leptoquark or leptoquarkino can be singly produced in such a collider. The scalar leptoquark will be produced in the association of an antiquark [16]. As discussed before in the usual SUSY theories, the leptoquark further decays into a quark and a positron with the final state consisting of positron +jets. The leptoquarkino however will be produced along with a squark. This leptoquarkino will then decay into a lepton and a squark. The final state has electron +jets+missingenergy. The missing energy part will then disentangle the leptoquarkino signal. In the gauge mediated SUSY breaking scenario, the final state has either a hard photon or $\tau^+\tau^-$ along with electron +jets+missing energy.

In e^+e^- collider the signal for this leptoquarkino would be jets +missing energy, since a squark anti-squark pair will be produced from a t channel exchange of leptoquarkino. The exchange of a t channel leptoquark however gives rise to jets without any missing energy. It is not possible for LEP II to see the signal of leptoquarkino due to the large squark mass. A higher energy machine is needed for that purpose. A detailed study of the discovery prospects of the leptoquarkino in different colliders will be presented elsewhere.

Another important implication of our model is that there does not seem to any operator which can give a non-negligible charged current signal at HERA. Thus observation of charged current like events above background at HERA will be an evidence against this model.

Let us now discuss the possible implications of the existence of a low mass leptoquark supermultiplet for unification. We will assume that the physics immediately beyond MSSM (i.e. in the TeV region) is described by a supersymmetric left-right symmetric model (SUSYLR) with the addition of the leptoquark multiplets described above. The evolution of the gauge couplings depend not only on the individual beta functions but also on the nature of the final unification group which determines the normalizations of the various gauge couplings at low energies. For instance, if we envision embedding the SUSYLR group within a simple group such as $SO(10)$ or $SU(16)$ etc, the properly normalized weak hypercharge, I_Y is given by the familiar formula $I_Y = \sqrt{\frac{3}{5}}(\frac{Y}{2})$ and the properly normalized $B - L$ charge I_{BL} is given by $I_{BL} = \sqrt{\frac{3}{2}}(\frac{B-L}{2})$. This leads to the matching formula for the weak hypercharge coupling to be $\alpha_{I_Y}^{-1} = \frac{3}{5}\alpha_{2R}^{-1} + \frac{2}{5}\alpha_{BL}^{-1}$ at M_R . Evolving our model with this kind of unification leads to a unification scale of about $10^{9.5} \text{ GeV}$ assuming that at the W_R scale, we have two bidoublets, right handed triplets of type $(1, 3, -2, 1) + (1, 3, +2, 1)$ without their left-handed partner in addition to the aforementioned leptoquark multiplets. This is therefore unacceptable since it will lead to catastrophic proton decay via the exchange of gauge bosons that violate baryon number in the $SO(10)$ models. We therefore consider a different kind of embedding of SUSYLR into an $SU(5) \times SU(5)$

model suggested in a recent paper [17].

SU(5) × SU(5) embedding of SUSYLR with leptoquarks

First let us briefly recapitulate the fermion assignments of the model: they belong to the $(\bar{\mathbf{5}} + \mathbf{10}, 1) + (1, \mathbf{5} + \bar{\mathbf{10}})$ multiplet [17] for every generation. One therefore needs a weak singlet vectorlike pair of D, U quarks and a heavy weak singlet vectorlike charged lepton $E^\pm$. The lefthanded $(\bar{\mathbf{5}}, 1)$ multiplet then consists of $(D_1^c, D_2^c, D_3^c, e^-, \nu_e)$ whereas the righthanded multiplet $(1, \mathbf{5})$ is given by $(D_1, D_2, D_3, e^c, \nu^c)$. The assignments to the $\mathbf{10}$ dimensional representations are easily obtained [17]. For instance, the fermion assignment in $(\mathbf{10}, 1)$ representation is given by

$$\begin{pmatrix} 0 & U_3^c & -U_2^c & u_1 & d_1 \\ -U_3^c & 0 & U_1^c & u_2 & d_2 \\ U_2^c & -U_1^c & 0 & u_3 & d_3 \\ -u_1 & -u_2 & -u_3 & 0 & E^+ \\ -d_1 & -d_2 & -d_3 & -E^+ & 0 \end{pmatrix} \quad (3)$$

and similarly for the $(1, \bar{\mathbf{10}})$ multiplet which contains the other chirality states for the above fields arranged exactly the same way. The Higgs mechanism of the model is implemented by multiplets of type $(\mathbf{5}, \bar{\mathbf{5}}) + (\bar{\mathbf{5}}, \mathbf{5})$ and they lead to the vectorlike quarks and leptons acquiring mass at the unification scale. This leaves the low energy theory to be the usual SUSYLR model. The SUSYLR symmetry is broken down to MSSM by the Higgs multiplets of type $(1, \mathbf{15}) + (\bar{\mathbf{15}}, 1)$ which also leads to the see-saw mechanism for neutrino masses. Below the W_R scale the theory is MSSM with the important difference that now the R-parity violating couplings are automatically absent. In order to accomodate the leptoquarks in this model, we include the multiplets $(\mathbf{5}, \mathbf{10}) + (\bar{\mathbf{10}}, \bar{\mathbf{5}})$. We assume that below the GUT scale, only members of these multiplets that remain light are the leptoquarks discussed above.

Turning to unification with leptoquarks in this model, we first note that since $U(1)_{B-L}$ and $SU(3)_c$ now emerge from two different $SU(5)$'s and that there are more fermions in the fundamental representation of the GUT group than the $SO(10)$ case, the normalization of the low energy couplings are totally different from the previous case. For instance, now $I_Y = \sqrt{\frac{3}{13}}(\frac{Y}{2})$ and $I_{BL} = \sqrt{\frac{3}{10}}(\frac{B-L}{2})$; furthermore, $I_a = \sqrt{\frac{1}{2}}(I_{aL} + I_{aR})$ where a denotes $SU(3)_c$ or $B - L$ generators. Because of this unification profile is now very different; however with the same field content at M_R as above, we find a unification scale of 10^9 GeV in the one loop approximation. In Table 1, we have showed some unification scenarios by varying the numbers of bidoublets. We have one $(2, 2, 4/3, 3) + (2, 2, -4/3, 3^*)$ and one $(1, 3, -2, 1) + (1, 3, 2, 1)$ and have used $\alpha = 1/127.9$, $\sin^2\theta_w = 0.2321$ and $\alpha_s = 0.118$ at

the M_Z scale. From the table I, one can see that one can have one unified coupling at the GUT scale using seven bidoublets. It is important to point out that this low unification scale does not conflict with proton lifetime since proton is absolutely stable in this model [17]. The essential reason for this is that the vectorlike quarks and the light quarks do not mix with each other due to the choice of the gauge group and Higgs representations.

We find it interesting that CDF collaboration, in their analysis of the dilepton channel, have reported [18] 4 events containing at least one τ (3 of the events have at least one b quark). Their estimated background from non- $t\bar{t}$ productions is 2 events. Using their measured cross-sections of 7.5 pb (from DIL, SVX and SLT channels), and their τ channel acceptance of 0.12%, only one such event is expected from the $t\bar{t}$ productions in the 110 pb $^{-1}$ data. It is entirely possible that this event is due to statistical fluctuation, or due to charged Higgs production, however such excess events are expected from the leptoquark or leptoquarkino pair productions. One leptoquark decays to a τ and the top quark, and the other to a e or a τ . A leptoquark (or leptoquarkino) pair production will have a branching ratio of about 0.75 for the dilepton channel with a τ in our model (compared to about 0.05 in $t\bar{t}$ productions). This will give an acceptance (branching ratio × detection efficiency) of about 1.8 %. Thus leptoquark (or leptoquarkino) pair production cross-section of about 0.5 pb will give rise to such an excess event. As can be seen from fig 1, this corresponds to produced leptoquarkino mass about 250 GeV (or a leptoquark of somewhat lower mass).

In conclusion, we have suggested an alternative leptoquark interpretation of the HERA high Q^2 anomaly within an extension of the MSSM in such a way that R-parity conservation is maintained automatically. The superpartner of the leptoquark, the leptoquarkino in this model leads to many interesting and testable predictions in present and future hadronic as well as in the future e-gamma colliders. At Tevatron, the pair productions of these leptoquarkinos give rise to very distinctive signal, namely opposite sign dilepton events of the type ee , $e\tau$, $\tau\tau$ accompanied by dijets plus missing energy, but no events of $\mu\mu$ type. If the leptoquarkino is not too much heavier than the “HERA leptoquark”, then these events will be observable in the present or upgraded Tevatron. It is possible that one excess event in the dilepton channel containing a τ , as reported by the CDF collaboration, could be due to the production of a leptoquark or leptoquarkino. Although clearly more data is needed before definitive conclusion can be drawn.

The work of B. D has been supported by a DOE grant no. DE-FG06-854ER-40224; the work of R. N. M. has been supported by NSF grant no. PHY-9421386 and the work of S. N. has been supported by DOE grant no. DE-FG02-94-ER40852.

- [1] C. Adloff et al., H1 collaboration, DESY 97-024, hep-ex/9702012.
- [2] J. Breitweg et al., ZEUS collaboration, DESY 97-025, hep-ex/9702015.
- [3] For early discussions of leptoquarks, see R. N. Mohapatra, G. Segre and L. Wolfenstein, Phys. Lett. **145B**, 433 (1984); W. Buchmuller, R. Ruckl and D. Wyler, Phys. Lett. **191B**, 442 (1987); S. Davidson, D. Bailey and B. A. Campbell, Z. Phys. **C61**, 613 (1994); M. Leurer, Phys. Rev. **D 50**, 536 (1994). J. L. Hewett, Proc. 1990 Summer Study on High Energy Physics, Snowmass, Co, ed. E.L. Berger, (world Scientific); J. Blumlein, hep-ph/9703287.
- [4] K. S. Babu, C. Kolda, J. March-Russell and F. Wilczek, hep-ph/9703299; M. C. Gozales-Garcia and M. Fabbri-chesi, hep-ph/9703346; G. Altarelli, J. Ellis, G. Giudice, S. Lola, M. Mangano, hep-ph/9703276; V. Barger et al., hep-ph/9703311; N. Di Bartolomeo and M. Fabbri-chesi, hep-ph/9703375; A. Nelson, hep-ph/9703379; W. Buchmuller and D. Wyler, hep-ph/9704317; K. Akama, K. Katsura and H. Terazawa, hep-ph/9704327.
- [5] S. Godfrey, hep-ph/9704380.
- [6] D. Choudhury and S. Raychaudhuri, hep-ph/9702392; G. Altarelli et al., Ref. 4; K. S. Babu et al., Ref. 4; H. Dreiner and P. Morawitz, hep-ph/9703279; J. Kalinowski et al., hep-ph/9703288; T. Kon and T. Kobayashi, hep-ph/9704221; J. Hewett and T. Rizzo, hep-ph/9703337; R. Barbieri, Z. Berezhiani and A. Strumia, hep-ph/9704275.
- [7] For a review see, H. Haber and G. Kane, Phys. Rep. **117**, 76 (1984).
- [8] For a review of the constraints on R-parity violating couplings, see G. Bhattacharyya, *Supersymmetry96*, edited by R. N. Mohapatra and A. Rasin, North Holland (1997), p. 83.
- [9] R. N. Mohapatra, Phys. Rev. **D 34**, 3457 (1986); M. Hirsch, H. Klapdor-Kleingrothaus and S. Kovalenko, Phys. Rev. Lett. **75**, 17 (1995).
- [10] R. N. Mohapatra, [9]; A. Font, L. Ibanez and F. Quevedo, Phys. Lett. **B 228**, 79 (1989); S. Martin, Phys. Rev. **D 46**, 2769 (1992).
- [11] R. M. Barnett et al. Phys. Rev. **D 54**, 1 (1996).
- [12] F. Abe et al., Phys. Rev. Lett. **74**, 2626 (1995); for a recent summary, see D. S. Kestelbaum, FERMILAB-CONF-97/016-E.
- [13] S. Abachi et al., Phys. Rev. Lett. **74**, 2632 (1995); N. Hadley, private communication for the most recent dilepton data.
- [14] C. Grosso-Pilcher (CDF-collaboration), talk at the Vanderbilt meeting, May 15th, 1997.
- [15] M. Dine and A. Nelson, Phys. Rev. **D47**, 1277 (1993); M. Dine, A. Nelson and Y. Shirman, Phys. Rev. **D51**, 1362 (1995); M. Dine, A. Nelson, Y. Nir and Y. Shirman, Phys. Rev. **D53**, 2658 (1996); M. Dine, Y. Nir and Y. Shirman, preprint SCIPP-96-30, hep-ph/9607397.
- [16] M. Doncheski, S. Godfrey, Talk given at 1996 DPF / DPB Summer Study on New Directions for High-Energy Physics (Snowmass 96), Snowmass, CO, 25 Jun - 12 Jul 1996, hep-ph/9612385 and references therein.
- [17] R. N. Mohapatra, Phys. Lett. **B379**, 115 (1996).
- [18] F. Abe et al (CDF collaboration), CDF/ANL/TOP/PUBLIC/4011,(1997).

$n_{\text{bidoublet}}$	$M_I(\text{GeV})$	$M_G(\text{GeV})$	$\alpha_{G_A}^{-1}$	$\alpha_{G_B}^{-1}$
5	10^3	$10^{9.11}$	5.12	2.29
6	10^3	$10^{8.84}$	4.05	3.46
7	$10^{3.75}$	$10^{9.04}$	4.25	4.25

TABLE II. The Grand unification scale, the intermediate scale and the strength of the unified couplings at the grand unified scale are shown for different numbers of bidoublets. α_{G_A} is the unified coupling for one $SU(5)$ group and α_{G_B} is the unified coupling for the other $SU(5)$ group




FIG. 1. Cross section for the production of leptoquarkino pair at Tevatron energy against the leptoquarkino mass.